

Proteotyping SARS-CoV-2 virus from nasopharyngeal swabs: a proof-of-concept focused on a 3 min mass spectrometry window

Duarte Gouveia^{1§}, Guylaine Miotello^{1§}, Fabrice Gallais¹, Jean-Charles Gaillard¹, Stéphanie Debroas¹, Laurent Bellanger¹, Jean-Philippe Lavigne^{2,3}, Albert Sotto⁴, Lucia Grenga¹, Olivier Pible¹, Jean Armengaud^{1*}

¹Université Paris Saclay, CEA, INRAE, Département Médicaments et Technologies pour la Santé (DMTS), SPI, 30200 Bagnols-sur-Cèze, France.

²U1047, Institut National de la Santé et de la Recherche Médicale, Université Montpellier, France

³VBMI, INSERM U1047, Université de Montpellier, Service de Microbiologie et Hygiène Hospitalière, CHU Nîmes, Nîmes, France.

⁴VBMI, INSERM U1047, Université de Montpellier, Service des Maladies Infectieuses et Tropicales, CHU Nîmes, Nîmes, France.

Supplementary data

Table S1. List of *m/z* values of the inclusion list.

Table S2. UNIPEPT taxonomical analysis of Swab R1, Swab R2, and simili swab 7R1-1pfu samples.

Table S3. List of peptide-to-spectrum matches (PSMs) assigned from the simili SARS-CoV-2 swabs (FDR<1%).

Table S4. List of proteins from the simili SARS-CoV-2 swabs and their spectral counts.

Table S5. List of peptide-to-spectrum matches (PSMs) assigned in the clinical samples (FDR<1%).

Table S6. List of proteins identified in the clinical samples and their spectral counts.

Table S7. Skyline report with experimental data on precursor ions from the nine swabs from COVID-19 patients.